arXiv:cond-mat/0312231v1 [cond-mat.stat-mech] 9 Dec 2003

Exact Solution for the Influence of Spectral Diffusion on Single-Molecule Photon-Statistics

Y. He, E. Barkai
Department of Chemistry and Biochemistry, Notre Dame University, Notre Dame, IN 46556.

(Dated: February 2, 2008)

We investigate the distribution of number of photons emitted by a single molecule undergoing a spectral diffusion process and interacting with a continuous wave field. Using a generating function formalism an exact analytical formula for Mandel's Q parameter is obtained. The solution exhibits transitions between: (i) Quantum sub-Poissonian and Classical super-Poissonian behaviors, and (ii) fast to slow modulation limits. Our solution yields the conditions on the magnitude of the spectral diffusion time scales on which these transitions are observed. We show how to choose the Rabi frequency in such a way that the Quantum sub-Poissonian nature of the emission process becomes strongest, we find $\frac{\Omega^2}{\Gamma} = \frac{\Gamma_{\mathrm{SD}}+\Gamma}{2}$ where Γ_{SD} (Γ) is the spectral diffusion (radiative) contribution to the width of the line shape.

PACS numbers: 82.37.-j, 05.10.Gg, 33.80.-b, 42.50.Ar

Physical, Chemical, and Biological systems are investigated in many laboratories using single molecule spectroscopy [1]. The investigation of the distribution of number of photons emitted from a single molecule source is the topic of extensive theoretical research e.g. [2, 3, 4, 5]. An important mechanism responsible for fluctuations in photon statistics from single molecule sources is spectral diffusion. In many cases the absorption frequency of the molecule will randomly change due to different types of interactions between the molecule and its environment (e.g. [1, 6, 7, 8] and Ref. therein). For example in low temperature glasses flipping two level systems embedded in the glassy environment, induce spectral jumps in the absorption frequency of the single molecule under investigation [6]. In this way the molecule may come in and out of resonance with the continuous wave laser field with which it is interacting.

Obviously a second mechanism responsible for fluctuations of photon counts is the quantum behavior of the spontaneous emission process [9, 10]. In his fundamental work Mandel [11] showed that a single atom in the process of resonance fluorescence, *in the absence of spectral diffusion*, exhibits sub-Poissonian photon statistics [12]. Photon statistics is characterized by Mandel's Q parameter

$$Q = \frac{\overline{N^2} - \overline{N}^2}{\overline{N}} - 1 \tag{1}$$

where N is the number of emitted photons within a certain time interval. The case $Q < 0$ is called sub-Poissonian behavior, while $Q > 0$ is called super-Poissonian. Sub-Poissonian statistics has no classical analog [11]. Briefly, the effect is related to anti-bunching of photons emitted from a single source and to Rabi-oscillations of the excited state population which favors an emission process with some periodicity in time. Sub-Poissonian statistics and anti-bunching were detected in several single molecule experiments, for example [13, 14].

In this Letter we obtain an exact analytical expression for the Q parameter in the long time limit, for a single molecule undergoing a stochastic spectral diffusion process. For that aim we use the powerful generating function method of Zheng and Brown [15]. If the spectral diffusion process is slow enough super-Poissonian behavior of Q is expected (see details in text). Our analytical expressions classify the transitions between sub and super Poissonian statistics. They give the conditions on the spectral diffusion time scale and magnitude of spectral jumps for sub-Poissonian non-classical behavior to be observed. Our result is valid for weak and strong excitation (i.e arbitrary Rabi frequency). It yields the lower bound on Q. The solution shows how in experiment we may choose the Rabi frequency so that the quantum nature of the photon emission process becomes larger, namely how to minimize Q in the Sub-Poissonian regime. This is important for the efficient detection of quantum effects in single molecule spectroscopy, since choosing too small or too large values of the Rabi frequency results in very small and hence undetectable values of Q. Finally our result is used to test the semi-classical linear response theory (i.e., weak Rabi frequency) of Barkai, Jung, and Silbey [7, 8]. The semi-classical theory yields $Q > 0$, while the main focus of this manuscript is on the quantum regime $Q < 0$.

Our starting point are the Zheng–Brown [15] generalized optical Bloch equations describing a chromophore with single excited and ground state

$$\begin{array}{l}
\dot{\mathcal{U}}\left(s\right) = -\frac{\Gamma}{2}\mathcal{U}\left(s\right) + \delta_L(t)\mathcal{V}\left(s\right)\\
\dot{\mathcal{V}}\left(s\right) = -\delta_L(t)\mathcal{U}\left(s\right) - \frac{\Gamma}{2}\mathcal{V}\left(s\right) - \Omega\mathcal{W}\left(s\right)\\
\dot{\mathcal{W}}\left(s\right) = \Omega\mathcal{V}\left(s\right) - \frac{\Gamma}{2}\left(1+s\right)\mathcal{W}\left(s\right) - \frac{\Gamma}{2}\left(1+s\right)\mathcal{Y}\left(s\right)\\
\dot{\mathcal{Y}}\left(s\right) = -\frac{\Gamma}{2}\left(1-s\right)\mathcal{W}\left(s\right) - \frac{\Gamma}{2}\left(1-s\right)\mathcal{Y}\left(s\right).
\end{array} \tag{2}$$

These equations are exact within the rotating wave approximation and optical Bloch equation formalism. In Eq. (2) Γ is the spontaneous emission rate of the electronic transition and Ω is the Rabi frequency. The time evolving detuning is $\delta_L(t) = \omega_L - \omega_0 - \Delta\omega(t)$, where ω_L (ω_0) is the laser frequency (the molecule's bare frequency), and $\Delta\omega(t)$ is the stochastic spectral diffusion process. The information about the photon statistics is contained in the moment generating function [15]

$$2\mathcal{Y}(s) \equiv \sum_{N=0}^{\infty} s^N P_N(t) \qquad (3)$$

where $P_N(t)$ is the probability of N emission events in the time interval $(0,t)$. Thus the solution of Eq. (2) yields the moments of N

$$\overline{N(t)} = 2\mathcal{Y}'(1) \qquad \overline{N^2(t)} = 2\mathcal{Y}''(1) + 2\mathcal{Y}'(1), \qquad (4)$$

with which the Mandel Q parameter can in principle be obtained. In Eq. (4), and in what follows, we use the notation $\frac{\partial}{\partial s} g(s)|_{s=1} \equiv g'(1)$ and similarly for second order derivatives with respect to s. The physical meaning of $\mathcal{U}(s)$, $\mathcal{V}(s)$, and $\mathcal{W}(s)$ and their relation to the standard Bloch equation was given in [15], and in closely related work by Mukamel [16], some discussion on this issue will follow Eq. (6). Note that when $s \to 1$ the damping terms in Eq. (2) become small [i.e. the $(1-s)\Gamma/2$ terms], hence relaxation of the generalized Bloch equations in the important limit of $s \to 1$ is slow.

In what follows we will consider the moments $\overline{N(t)}$, $\overline{N^2(t)}$. For this aim it is useful to derive equations of motion for the vector $z = \{\mathcal{U}(1), \mathcal{V}(1), \mathcal{W}(1), \mathcal{Y}(1), \mathcal{U}'(1), \mathcal{V}'(1), \mathcal{W}'(1), \mathcal{Y}'(1), \mathcal{Y}''(1)\}$. Taking the first and the second derivative of Eq. (2) with respect to s and setting $s = 1$, we find

$$\dot{z} = M(t) z \qquad (5)$$

where $M(t)$ is a 9×9 matrix $M(t) =$

$$\begin{pmatrix}
-\frac{\Gamma}{2} & \delta_L(t) & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
-\delta_L(t) & -\frac{\Gamma}{2} & -\Omega & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & \Omega & -\Gamma & -\Gamma & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & -\frac{\Gamma}{2} & \delta_L(t) & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & -\delta_L(t) & -\frac{\Gamma}{2} & -\Omega & 0 & 0 \\
0 & 0 & -\frac{\Gamma}{2} & -\frac{\Gamma}{2} & 0 & \Omega & -\Gamma & -\Gamma & 0 \\
0 & 0 & \frac{\Gamma}{2} & \frac{\Gamma}{2} & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & \Gamma & \Gamma & 0
\end{pmatrix} \qquad (6)$$

The first three lines of $M(t)$ describe the evolution of $\mathcal{U}(1), \mathcal{V}(1), \mathcal{W}(1)$, these are the standard optical Bloch equations in the rotating wave approximation. The fourth line of $M(t)$ is zero, it yields $\dot{\mathcal{Y}}(1) = 0$, this equation describes the normalization condition of the problem namely $\mathcal{Y}(1) = 1/2$ for all times t [to see this use Eq. (3) and $\sum_{N=0}^{\infty} P_N(t) = 1$]. The evolution of the remaining terms $\mathcal{U}'(1), \mathcal{V}'(1), \mathcal{W}'(1), \mathcal{Y}'(1), \mathcal{Y}''(1)$ are of current interest since they describe the fluctuation of the photon emission process. Solutions of time dependent equations like Eq. (5) are generally extremely difficult to obtain, a formal solution is given in terms of the time ordering operator T, $z(t) = T \exp[\int_0^t M(t) \mathrm{d}t] z(0)$.

Eq. (5) yields a general method for the calculation of Q for a single molecule undergoing a spectral diffusion process. The aim of this Letter is to obtain an exact solution of the problem for an important stochastic process used by Kubo and Anderson to investigate characteristic behaviors of line shapes. We assume $\Delta\omega(t) = \nu h(t)$ where ν describe frequency shifts, and $h(t)$ describes a random telegraph process: $h(t) = 1$ or $h(t) = -1$. The transition rate between state up (+) and state down (-) and vice versa is R. This stochastic process was used to describe behavior of line shapes in many systems [6, 20], here our aim is to calculate Q describing the line shape fluctuations.



FIG. 1: In (a) we show the line shape $I(\omega_L)$ vs ω_L for $\Gamma = 40$ MHz, $\Omega = \Gamma/\sqrt{2}$, and $\nu = 2\Gamma$. The line exhibits the well known transition between fast ($R = 100\Gamma$ dotted solid curve) and slow modulation limits ($R = \Gamma/10$ solid curve). In (b) we show the Q parameter, using the same parameters. Three type of behaviors are observed: (i) for fast modulation the Q parameter attains its minimum on zero detuning. (ii) For an intermediate modulation, $R = \Gamma/2$ (dotted curve) we obtain $Q < 0$ however now the minimum of Q is on $\omega_L - \omega_0 = \pm\nu$. (iii) For the slow process Q exhibits super-Poissonian statistics $Q > 0$. Note that we used $\Omega = \Gamma/\sqrt{2}$ in order to maximize the absolute value of Q in the fast modulation limit.

We use Burshtein's method [17, 18] of marginal averages, to solve the stochastic differential matrix equation (5). The method yields the average $\langle z \rangle$ with respect to the stochastic process. We will calculate $\langle z \rangle$ in the limit of long times, and then obtain the steady state behavior of the line shape $I(\omega_L) = \lim_{t\to\infty} \frac{\mathrm{d}}{\mathrm{d}t} \langle \overline{N(t)} \rangle$ and the Mandel Q parameter. Let $\langle z\rangle_\pm$ be the average of $z(t)$ under the condition that at time t the value of $h(t)=\pm 1$ respectively. $\langle z\rangle_\pm$ are called marginal averages, the complete average is $\langle z\rangle = \langle z\rangle_+ + \langle z\rangle_-$. The equation of motion for the marginal averages is

$$\begin{pmatrix} \langle \dot{z}_+\rangle \\ \langle \dot{z}_-\rangle \end{pmatrix} = \begin{pmatrix} M_+ - RI & RI \\ RI & M_- - RI \end{pmatrix} \begin{pmatrix} \langle z_+\rangle \\ \langle z_-\rangle \end{pmatrix}. \qquad (7)$$

In Eq. (7) the matrix $M_\pm$ is identical to matrix M in Eq. (6) when $\delta_L(t)$ is replaced by $\delta_L^\pm = \omega_L - \omega_0 \mp \nu$, and I is a 9×9 identity matrix. The 18×18 matrix in Eq. (7) is traceless, hence the time dependent solution of Eq. (7) is not obtained by direct inversion of it. We found the long time behavior of Eq. (7) after a long journey which involved the following five steps. **1**. We obtained the steady state solution for $\langle\{\mathcal{U}(1)_\pm, \mathcal{V}(1)_\pm, \mathcal{W}(1)_\pm, \mathcal{Y}(1)_\pm, \mathcal{Y}'(1)_\pm\}\rangle$. **2**. We used the solutions obtained in previous step to obtain inhomogeneous equations for $x = \langle\{\mathcal{U}'(1)_+, \mathcal{V}'(1)_+, \mathcal{W}(1)_+, \mathcal{U}'(1)_-, \mathcal{V}'(1)_-, \mathcal{W}'(1)_-\}\rangle$

$$\dot{x} = Ax + b(t) \qquad (8)$$

where

$$A = \begin{pmatrix} -\frac{\Gamma}{2} - R & \delta_L^+ & 0 & R & 0 & 0 \\ -\delta_L^+ & -\frac{\Gamma}{2} - R & -\Omega & 0 & R & 0 \\ 0 & \Omega & -\Gamma - R & 0 & 0 & R \\ R & 0 & 0 & -\frac{\Gamma}{2} - R & \delta_L^- & 0 \\ 0 & R & 0 & -\delta_L^- & -\frac{\Gamma}{2} - R & -\Omega \\ 0 & 0 & R & 0 & \Omega & -\Gamma - R \end{pmatrix} \qquad (9)$$

and $b(t) = (0, 0, b_+(t), 0, 0, b_-(t))$ with

$$b_\pm(t) = -\frac{\Gamma}{8} - \frac{\Gamma}{2}\mathcal{W}^{ss}(1)_\pm - \frac{\Gamma^2 t}{8}\left\{1 + 2\left[\mathcal{W}^{ss}(1)_+ + \mathcal{W}^{ss}(1)_-\right]\right\} \pm \frac{\Gamma^2}{8R}\left[\mathcal{W}^{ss}(1)_- - \mathcal{W}^{ss}(1)_+\right]. \qquad (10)$$

and

$$\mathcal{W}^{ss}(1)_+ = \frac{\Gamma}{4}\left[\left(A^{-1}\right)_{33} + \left(A^{-1}\right)_{36}\right] \qquad \mathcal{W}^{ss}(1)_- = \frac{\Gamma}{4}\left[\left(A^{-1}\right)_{63} + \left(A^{-1}\right)_{66}\right]. \qquad (11)$$

We note that $\mathcal{W}^{ss}(1)_\pm$ yields the steady state (ss) marginal averages of the population difference between the upper excited state and ground state. **3**. Using Mathematica we found an analytical expression for A^{-1}, the inverse of the Matrix A, in terms of the parameters of the problem $R, \nu, \Gamma, \omega_L - \omega_0, \Omega$. **4**. Using A^{-1} and Eq. (8) we find the long time behavior of the vector x. We then use Eq. (7) to obtain steady state behavior of $\langle\mathcal{Y}''(1)^{\mathrm{ss}}\rangle_\pm$. **5**. Finally we obtain the Q parameter using: $Q = \left[\langle\mathcal{Y}''(1)^{\mathrm{ss}}\rangle - 2\langle\mathcal{Y}'(1)^{\mathrm{ss}}\rangle^2\right]/\langle\mathcal{Y}'(1)^{\mathrm{ss}}\rangle$.

Following these steps we obtain the main result of this Letter

$$Q = \frac{\Gamma^2}{2}\sum_{i=3,6}\sum_{j=3,6}\left(A^{-1}A^{-1}\right)_{ij} + \frac{\Gamma^2}{RI(\omega_L)}\left[\mathcal{W}^{ss}(1)_+ - \mathcal{W}^{ss}(1)_-\right]^2 + \frac{\Gamma}{I\left(\omega_L\right)}\sum_{k=\pm}\left[\mathcal{W}^{ss}(1)_k\left(1 + 4\mathcal{W}^{ss}(1)_k\right)\right], \qquad (12)$$

where the line is

$$I\left(\omega_L\right) = \Gamma\left[\frac{1}{2} + \mathcal{W}^{ss}(1)_+ + \mathcal{W}^{ss}(1)_-\right]. \qquad (13)$$

The general expression for Q in terms of original parameters of the problem can be found in [19]. Below we will analyze the physical behaviors of Q obtaining simple equations in limiting cases.

In Fig. 1 we show the line shape and the Q parameter. We choose $\Omega = \Gamma/\sqrt{2}$, fix ν, and vary the rate R. The line exhibits two well known behaviors: a fast modulation limit where the line is Lorentzian and a slow modulation limit where the line exhibits two peaks on $\omega_L = \pm\nu$. The Q parameter exhibits three behaviors: (i) fast modulation limit, where the single molecule exhibits a sub-Poissonian behavior, and Q attains its minimum on zero detuning. (ii) Intermediate modulation limit where photon statistics is still sub-Poissonian, however now Q attains its minimum on $\omega_L = \pm\nu$. And (iii) a slow modulation limit where a super-Poissonian behavior is observed. We now analyze these limits in some detail. Without loss of generality we use units where $\Gamma = 1$, and set $\omega_0 = 0$ hence ω_L is the detuning frequency.

For zero detuning, $\omega_L = 0$, we obtain from Eq. (12)

$$Q = -\frac{2\,\Omega^2\left[4\,\nu^2\,\left(-1 + 4\,R\right) + 3\,\left(1 + 4\,R\right)^2\right]}{\left[4\,\nu^2 + \left(1 + 2\Omega^2\right)\,\left(1 + 4\,R\right)\right]^2}. \qquad (14)$$

Eq. (14) indicates a transition from sub-Poissonian statistics ($Q < 0$) to super-Poissonian statistics ($Q > 0$) when $4\nu^2 = 3(1+4R)^2/(1-4R)$. For fast processes satisfying $R > 1/4$, (i.e. $R > \Gamma/4$ in natural units) we find sub-Poissonian behavior for all values of ν and Ω. As mentioned in introduction an important issue is how to choose Ω in such a way to minimize Q in the sub-Poissonian regime (i.e., how to maximize the quantum nature of the photon statistics). Using Eq. (14) it is easy to find the lower bound $Q \geq -3/4$. The absolute minimum $Q = -3/4$ is found when $\Omega = 1/\sqrt{2}$ and $\nu = 0$, namely for a stable molecule whose absorption frequency is not fluctuating. For fixed ν and R the largest quantum fluctuations are observed when $\Omega^2_{\text{min}} = (1 + 4\nu^2 + 4R)/[2(1 + 4R)]$. Note however that for fast processes $\nu << R$, the parameters ν and R cannot be directly deduced in experiment, due to motional narrowing effect. In Eq. (17) we will give an expression for Ω_{min} in terms of physical observables.

Consider the fast modulation limit. If we take $R \to \infty$ keeping Ω and ν fixed we obtain from Eq. (12)

$$\lim_{R\to\infty} Q = -\frac{2\Omega^2\left(3 - 4\omega_L^2\right)}{\left(1 + 2\Omega^2 + 4\omega_L^2\right)^2}. \tag{15}$$

This result was derived by Mandel for an atomic transition in the *absence* of spectral transition. As expected when the time scale for spectral jumps becomes very short, the molecule does not respond to the stochastic fluctuation. A more physically interesting case is to let $R \to \infty$ and $\nu \to \infty$ but keep the spectral diffusion (SD) contribution to the line width $\Gamma_{\text{SD}} \equiv \nu^2/R$ remain finite. In this limit, the line shape is Lorentzian and it exhibits motional narrowing (i.e. as R is increased the line becomes narrower). In this fast limit we obtain from Eq. (12)

$$Q_{\text{fast}} = -\frac{2\Omega^2\left[3 + 5\Gamma_{\text{SD}}^2 + \Gamma_{\text{SD}}^3 - 4\omega_L{}^2 + \Gamma_{\text{SD}}\left(7 + 4\omega_L^2\right)\right]}{\left[1 + \Gamma_{\text{SD}}{}^2 + 2\Omega^2 + 2\Gamma_{\text{SD}}\left(1 + \Omega^2\right) + 4\omega_L^2\right]^2}. \tag{16}$$

Note that photon statistics are sub-Poissonian provided that the detuning is not too large. The Q parameter obtains its minimum when

$$\Omega^2_{\text{min}} = \frac{\left(1 + \Gamma_{\text{SD}}\right)^2 + 4\omega_L^2}{2\left(1 + \Gamma_{\text{SD}}\right)}, \tag{17}$$

and for zero detuning $Q_{\text{min}} = -(\Gamma_{\text{SD}} + 3)/4(1 + \Gamma_{\text{SD}})$. The simple Eq. (17) is important, since it shows how to choose the Rabi frequency in order to obtain strong sub-Poissonian behavior. Since Γ_{SD} is a physical observable (unlike R and ν) Eq. (17) is of practical value.

To investigate the intermediate and slow modulation limit we consider the value of Q for $\omega_L = \nu$. We consider a strong coupling limit $\nu >> 1$, (e.g. spectral jumps of the order of $\nu =$ GHz $>> \Gamma$). From Eq. (12) we obtain

$$\lim_{\nu\to\infty} Q_{\omega_L=\nu} =$$

$$-\frac{\left(1 + 2R\right)\Omega^2\left[-1 + 16R^2 + 8R^3 - 2\Omega^2 + 2R\left(2 + \Omega^2\right)\right]}{2R\left[1 + 4R^2 + 2\Omega^2 + 2R\left(2 + \Omega^2\right)\right]^2}. \tag{18}$$

It is easy to see that Eq. (18) exhibits both sub-Poissonian and super–Poissonian behaviors. When the process is very *slow*, namely $R \to 0$, we obtain

$$\lim_{\nu\to\infty} Q_{\omega_L=\nu} \sim \frac{\Omega^2}{2R\left(1 + 2\Omega^2\right)} \tag{19}$$

a super-Poissonian behavior. In the *intermediate modulation limit*, for example when $R = 1$, we obtain

$$\lim_{\nu\to\infty} Q_{\omega_L=\nu} = -\frac{81\Omega^2}{2\left(9 + 4\Omega^2\right)^2} \tag{20}$$

a sub-Poissonian behavior. In this case the fluctuations are strongest when $\Omega^2_{\text{min}} = 9/4$ and then $Q_{\text{min}} = -9/32$. It is important to notice that in the same limit $\nu \to \infty$ we have for zero detuning $|Q| \propto \nu^{-2} \to 0$ hence in the slow and intermediate modulation limits the fluctuations on $\omega_L = \pm\nu$ are much stronger than the fluctuations on zero detuning (an example is in Fig. 1).

The detailed interaction between a chromophore and a bath is usually extremely complex, still for many fast processes line shapes are simple Lorentzians (universality). We expect that certain behaviors of Q obtained in this work are also universal, mainly the behavior in the fast modulation limit and the transition between fast, intermediate, and slow limits. We also expect, but have no proof, that the maximum of the quantum fluctuations, is attained close to $\Omega^2_{\text{min}}/\Gamma = (\Gamma_{\text{SD}} + \Gamma)/2$ [i.e., Eq. (17) in natural units and for zero detuning], also in other models of line broadening. Finally, the optimization of external fields to obtain strong sub-Poissonian and anti-bunching signals from single molecules is a new topic of research worthy of further investigation.

---

[1] W. E. Moerner and M. Orrit, *Science* **283** 1670 (1999).

[2] D. E. Makarov, H. Metiu *J. Chem. Phys.* **115** 5989–5993 (2001)

[3] G. K. Schenter, H. P. Lu, X. S. Xie *J. Phys. Chem. A* **103** 10477 (1999).

[4] A. M. Berezhkovskii, A. Szabo, and G. H. Weiss *J. Phys. Chem. B* **104** 3776 (2000).

[5] V. Barsegov, S. Mukamel *J. of Chem. Phys.* **116** 9802 (2002).

[6] E. Geva, and J. L. Skinner, *J. Phys. Chem. B* **101**, 8920 (1997).

[7] E. Barkai, Y. Jung, R. Silbey *Phys. Rev. Lett.* **87** 207403 (2001)

[8] Y. Jung, E. Barkai, and R. Silbey *Adv. Chem. Physics* **123**, 199 (2002) and cond-mat/0311428

[9] L. Mandel, and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, New York, 1995).

[10] M. B. Plenio, and P. L. Knight *Rev. of Mod. Phys.* 70:101–144 (1998).

[11] L. Mandel, *Opt. Lett.* **4** 205 (1979).

[12] R. Short, and L. Mandel, *Phys. Rev. Lett.* **51** 384 (1983).

[13] Th. Basche, W. E. Moerner, M. Orrit, and H. Talon,

Phys. Rev. Lett. **69**, 1516 (1992).

[14] L. Fleury, J-M. Segura, G. Zumofen, B. Hecht, and U. P. Wild, *Phys. Rev. Lett.* **84**, 1148 (2000).

[15] Y. Zheng, F. L. H. Brown *Phys. Rev. Lett.* **90** 238305 (2003)

[16] S. Mukamel, cond-mat/0307768

[17] A. I. Burshtein *Sov. Phys. JETP* **22** 937 (1966).

[18] B. W. Shore *The Theory of Coherent Atomic Excitation* Vol. 2, Wiley New-York (1990).

[19] We suggest that exact expression for Q will be published as an EPAPS document, see http://www.aip.org./pubservs/epaps.html

[20] B. J. Berne, and R. Pecora *Dynamic Light Scattering* Dover Publication, Chapter 6.

I. EXACT SOLUTION

The exact solution for the Q parameter, with $\Gamma = 1$, in the long time limit, $Q = \text{Numerator}[Q]/\text{Denominator}[Q]$ is

$$\text{Denominator}[Q] =$$

$$\begin{aligned}
&R\,(4{\omega_L}^2\,(1+2\,R)+4\,\nu^2\,(1+6\,R+8\,R^2)+(1+4\,R)\,(1+6\,R+8\,R^2+2\,\Omega^2))\,(16\,\nu^4\,(1+2\,R)+\\
&8\,\nu^2\,(1-4\,{\omega_L}^2\,(1+2\,R)+2\,\Omega^2+2\,R\,(3+4R)\,(1+\Omega^2))+(1+4\,{\omega_L}^2+2\,\Omega^2)\\
&(4{\omega_L}^2\,(1+2\,R)+(1+4\,R)\,(1+6\,R+8\,R^2+2\,\Omega^2)))^2
\end{aligned} \tag{21}$$

$$\text{Numerator}[Q] =$$

$$\begin{aligned}
&-2\,\Omega^2\,\Big(256\,\nu^8\,R\,(1+2\,R)^3\,(-1+16\,R^2)-(-3+4\,\omega_L^2)\,R\,({\omega_L}^2\,(4+8\,R)+(1+4\,R)\,(1+6\,R+8\,R^2+2\,\Omega^2))^3\\
&-128\,\nu^6\,(1+2\,R)^2\,(4\,{\omega_L}^2\,(1+4\,R+4\,R^2+16\,R^3+32\,R^4)-3\,R\,(1+4\,R)\,(24\,R^2+32\,R^3-\Omega^2+4\,R\,(1+\Omega^2)))\\
&+32\,\nu^4\,(1+2\,R)\,\Big(16\,{\omega_L}^4\,(1+2\,R)^2\,(2+5\,R+8\,R^3)+4\,{\omega_L}^2\,(1+6\,R+8\,R^2)\,(-2+112\,R^4+64\,R^5-4\,\Omega^2+\\
&R^2\,(76-8\,\Omega^2)-R\,(-4+\Omega^2)+16\,R^3\,(10+\Omega^2))+3\,R\,(1+4\,R)\,(1+832\,R^4+512\,R^5+\Omega^2-2\,\Omega^4+32\,R^3\,(16+5\,\Omega^2)+\\
&4\,R^2\,(37+32\,\Omega^2)+R\,(20+26\,\Omega^2+8\,\Omega^4)))-8\,\nu^2\,\Big(64\,{\omega_L}^6\,(1+2\,R)^4-R\,(1+4\,R)\,(1+6\,R+8\,R^2+2\,\Omega^2)^2\,(4+152\,R^2\\
&+160\,R^3-\Omega^2+4\,R\,(11+\Omega^2))-16\,{\omega_L}^4\,(1+2\,R)^2\,(-2+40\,R^3-4\,\Omega^2-4\,R^2\,(-5+6\,\Omega^2)-R\,(4+17\,\Omega^2))-4\,{\omega_L}^2\\
&(1+2\,R)\,\Big(7680\,R^6+2048\,R^7+256\,R^5\,(37+\Omega^2)-(1+2\,\Omega^2)^2+32\,R^4\,(169+36\,\Omega^2)+8R^3\,(191+108\,\Omega^2-8\,\Omega^4)+\\
&4R^2\,(47+43\,\Omega^2-12\,\Omega^4)-2\,R\,(-1+5\,\Omega^2+14\,\Omega^4)\Big)\Big)\Big)\Big)\,.
\end{aligned} \tag{22}$$

The line shape ($\Gamma = 1$) is

$$I\,(\omega_L) =$$

$$\begin{aligned}
&[\Omega^2\,(1+32\,R^2+32\,R^3+4\,\nu^2\,(1+6\,R+8\,R^2)+2\,\Omega^2+4\,\omega_L^2+2\,R\,(5+4\,\Omega^2+4\,\omega_L^2))]\,/\\
&[16\,\nu^4\,(1+2\,R)+8\,\nu^2\,(1+2\,\Omega^2+8\,R^2\,(1+\Omega^2)-4\,\omega_L^2+R\,(6+6\,\Omega^2-8\,\omega_L^2))+\\
&(1+2\,\Omega^2+4\,\omega_L^2)\,(1+32\,R^2+32\,R^3+2\,\Omega^2+4\,\omega_L^2+2\,R\,(5+4\,\Omega^2+4\,\omega_L^2))]
\end{aligned} \tag{23}$$